



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let $\cos\beta\sin\theta=\sin\varphi$; then the second term becomes

$$2a^3 \int_0^{\frac{1}{2}\pi-\beta} \sin^4\varphi \cos^2\varphi \log \left[\frac{\cos\varphi \cos\beta + \sqrt{(\cos^2\beta - \sin^2\varphi)}}{\sin\beta \sin\varphi} \right] d\varphi = 2a^3 \int_0^{\frac{1}{2}\pi-\beta} \left(\frac{1}{16}\varphi - \frac{1}{16}\sin\varphi \cos\varphi - \frac{1}{4}\sin^3\varphi \cos\varphi + \frac{1}{6}\sin^5\varphi \cos\varphi \right) \frac{\cos\beta d\varphi}{\sin\varphi \sqrt{(\cos^2\beta - \sin^2\varphi)}}.$$

$$\therefore V = \frac{1}{16}\pi a^3 \cos^3\beta \cot^2\beta (1 + \sin^2\beta) + \frac{1}{8}a^3 \cos\beta \int_0^{\frac{1}{2}\pi} d\theta + \frac{1}{2}\cos^3\beta \int_0^{\frac{1}{2}\pi} \sin^2\theta d\theta - \frac{1}{8}a^3 \cos^5\beta \int_0^{\frac{1}{2}\pi} \sin^4\theta d\theta - \frac{1}{8}a^3 \int_0^{\frac{1}{2}\pi} \frac{\sin^{-1}(\cos\beta \sin\theta) d\theta}{\sin\theta \sqrt{(1 - \cos^2\beta \sin^2\theta)}}.$$

$$\therefore V = \frac{1}{16}\pi a^3 \cot\beta \operatorname{cosec}\beta - \frac{1}{4}\pi a^3 \cos^3\beta$$

$$- \frac{1}{8}a^3 \int_0^{\frac{1}{2}\pi} \frac{\sin^{-1}(\cos\beta \sin\theta) d\theta}{\sin\theta \sqrt{(1 - \cos^2\beta \sin^2\theta)}} - \frac{1}{8}a^3 \int_0^{\frac{1}{2}\pi} \frac{\sin^{-1}(\cos\beta \sin\theta) d\theta}{\sin\theta \sqrt{(1 - \cos^2\beta \sin^2\theta)}}$$

$$= \frac{1}{8}a^3 \cos\beta \int_0^{\frac{1}{2}\pi} (1 + \frac{2}{3}\cos^2\beta \sin^2\theta + \frac{8}{15}\cos^4\beta \sin^4\theta + \frac{16}{35}\cos^6\beta \sin^6\theta + \dots) d\theta$$

$$= \frac{1}{16}\pi a^3 (\cos\beta + \frac{1}{3}\cos^3\beta + \frac{1}{5}\cos^5\beta + \frac{1}{7}\cos^7\beta + \dots) = \frac{1}{16}\pi a^3 \log \cot \frac{1}{2}\beta.$$

$$\therefore V = \frac{1}{4}\pi a^3 (3\cot\beta \operatorname{cosec}\beta - 2\cos^3\beta - 3\log \cot \frac{1}{2}\beta).$$


150. Proposed by E. B. ESCOTT, Instructor in Mathematics, University of Michigan, Ann Arbor, Mich.

Find total area between the curve $x^4y - x^2 + 4y - 1 = 0$ and the x -axis.

Solution by J. E. SANDERS, Hackney, Ohio, and the PROPOSER.

The equation may be written $y = \frac{x^2 + 1}{x^4 + 4}$.

$$\text{Area} = 2 \int_0^\infty y dx = 2 \int_0^\infty \frac{x^2 + 1}{x^4 + 4} dx$$



$$= \frac{1}{4} \left[\int_0^\infty \frac{x+2}{x^2-2x+2} dx - \int_0^\infty \frac{x-2}{x^2+2x+2} dx \right] = \frac{1}{8} \left[\log(x^2-2x+2) \right.$$

$$\left. - \log(x^2+2x+2) + 6[\tan^{-1}(x+1) + \tan^{-1}(x-1)] \right]_0^\infty$$

$$= \frac{1}{8} \left[\log \frac{x^2-2x+2}{x^2+2x+2} + 6 \tan^{-1} \frac{2x}{2-x^2} \right]_0^\infty = \frac{1}{8}(6\pi) = \frac{3}{4}\pi. \quad \text{Answer.}$$

$\left[\tan^{-1} \frac{2x}{2-x^2} \right]_{x=\infty} = \tan^{-1} 0 = \pi$, since the integrand has increased and passed through ∞ for $x=1/\sqrt{2}$.

Also solved by G. B. M. ZERR.

MECHANICS.

144. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Pressure is applied perpendicularly to the plane surface yz , bounding an otherwise infinite isotropic solid. Find the resultant displacements, if the pressure varies as $\sin\left(\frac{2\pi y}{a}\right) + \sin h\left(\frac{2\pi y}{a}\right)$.

No solution of this problem has been received.

145. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

$ABCD$, $GCEF$ are equal parallelograms, DCG and BCE being straight lines. If the figure be considered as formed of smooth light jointed bars and if BD be a light rod, and the whole be suspended from A , find the stress in BD if a weight be hung from F . Also find the stress if a light rod GE replace BD .

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Since the bars are light we can disregard their weight. Let P be the weight. Then by virtual work

$$Pd(AC) + Sd(BD) = 0 \dots (1).$$

$$\text{But } AC^2 + BD^2 = 2AD^2 + 2DC^2.$$


$$\therefore ACd(AC) + BDd(BD) = 0 \dots (2).$$

From (1) and (2),

$$\frac{d(AC)}{d(BD)} = -\frac{S}{P} = -\frac{BD}{AC}.$$

$$\therefore S = \frac{P \cdot BD}{AC}. \quad \text{Similarly for } GE.$$

$$S_1 = \frac{P \cdot GE}{CF}. \quad \text{The stress is the same for both.}$$



146. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

A diffraction grating, with lines .05 mm. apart is held in front of a Bunsen's burner in which common salt is volatilized, and, when viewed through a telescope it is found that the angular distances of the first, second, third, fourth, fifth, and sixth bright bands from the central one are respectively $41'$, $1^\circ 21'$, $2^\circ 2'$, $2^\circ 42'$, $3^\circ 23'$ and $4^\circ 3'$. Required the wave length of sodium light.